

# A QUON MODEL

ZHENGWEI LIU, ALEX WOZNIAKOWSKI, AND ARTHUR JAFFE  
HARVARD UNIVERSITY, CAMBRIDGE, MA 02138

ABSTRACT. We present a new, three-dimensional, topological model for quantum information. Our picture combines charged excitations carried by strings, with topological properties that arise from embedding the strings in a three manifold. A quon is a neutral pair of strings. The properties of the manifold allow us to interpret multi-quons and their transformations in a natural way. We obtain a new type of relation, a string-genus “joint relation,” between (1): a diagram that contains a neutral string surrounding the genus of the manifold, and (2): the diagram without the string and genus. We use the joint relation to obtain a 3D topological interpretation of the  $C^*$  Hopf algebra relations, that are widely used in tensor networks. We obtain a 3D representation of the CNOT gate and a 3D topological protocol for teleportation.

## 1. INTRODUCTION

Topological quantum information was formulated by Kitaev [1] and Freedman et al. [2]. Here we study a new, topological model for quantum information that we call the *quon* model, where the name quon suggests a quantum particle.

In earlier work we found a *four-string model*, to represent a 1-qudit vector as a neutral particle-antiparticle pair [3, 4]. Our four-string model of a qudit provides one example of a quon. A central feature of our picture is that the quon is a *neutral* pair, even though it is composed of strings that may be charged. For fermions neutrality of a pair is automatic, as any two fermions are bosonic (the condition for multi-fermion neutrality). But excitations with more than two charge states, such as parafermions, require pairing an excitation with its anti-excitation, to obtain a neutral composite.

The neutrality of pairs makes finding a corresponding multi-quon picture difficult. Braiding strings from two different quons can destroy neutrality. So can one characterize transformations that map the multi-quon space into itself? Here we solve this problem and give a full, multi-quon model. Our construction provides a general four-string quon structure, of which multi-qudits are a special case.

Our new idea is to consider the quon strings as embedded into a topological three-manifold. Thus the quon model has the flavor of a topological field theory with additional relations that involve both the manifold and the strings. We call these *joint relations*. In §9–11 we use these joint relations to obtain a natural understanding of Frobenius and  $C^*$ -Hopf algebra relations. An important such relation in our model allows one to remove a genus from the manifold, when that genus is surrounded by a neutral string.


1.1. **1-Quons.** Our diagram for a 1-quon is a hemispherical cap with no input points and four output points, namely a diagram of the form:



The dotted box designates the internal structure that specifies the quon vector. In case a quon is a qudit of degree  $d$ , one has a simple representation for a 1-quon basis: the interior of a cap contains two charged strings,

each linking two of the output points. The value of the charge on one sting may equal either  $0, 1, \dots, d-1 \in \mathbb{Z}_d$ , while the other string carries the negative of that charge. For  $d = 2$  the quons reduce to fermions. The four-string model for a qudit found in §5.3 of [3] arose as a natural generalization of Kitaev's picture [5] of a spin as a pair of fermions. The four strings arise as we represent the Pauli matrices  $X, Y, Z$  by four parafermions. In our reinterpretation, we replace the two fermions by a pair of parafermion/anti-parafermion unitaries with opposite charge. We represent transformations on 1-quons as a box with four input points and four output points, embedded in a three-manifold. We describe various bases in §4.

**1.2. Multi-quons.** For multi-quons, one has a hemisphere cap for each 1-quon. The manifold for a transformation on  $n$ -quons has  $n$  input handles to attach to the  $n$  caps, with strings attaching as well. This leads to a natural multi-particle structure, and it allows us to analyze the full Hilbert space for multi-quons. Each individual quon remains a neutral pair. For example, a three-quon to three-quon transformation has the form:



In our previous work we represented qudits with a two-string model. In this case the picture of multi-qudits is more straightforward, and fits into the framework of planar para algebras [3, 4, 6, 7]. Para-isotopy relations play an important role, as they reflect the parafermion multiplication law. In the quon model, para isotopy also plays a role, but the neutrality condition for quons means that quons satisfy isotopy. In the quon model, we also need the joint topological relations between strings and the ambient manifold which carries the strings.

**1.3. Quons as topological algebra.** Abramsky and Coecke studied quantum information from a categorical point of view [8]. The Hopf algebra axioms have been studied in tensor networks and lead to applications in quantum information [9, 10, 11, 12, 13, 14]. Vicary and Reutter have studied other concepts from a higher category and from a planar algebra point of view [15, 16].

In our quon we represent pictures in 3D space. This captures the categorical structures in two directions. In particular, we obtain Frobenius algebras in the  $X$ -direction and the  $Y$ -direction corresponding to the maps COPY and SUM in tensor networks. They define the underlying Hilbert space as a  $C^*$ -Hopf algebra. Moreover, the string Fourier transform  $\mathfrak{F}_S$  is a  $90^\circ$  rotation around the  $Z$ -axis. Conjugation by  $\mathfrak{F}_S$  maps one Frobenius algebra to the other. This gives a topological interpretation of the algebraic axioms of Hopf algebras.

The defining algebraic relations reduce to the invariance of certain elementary diagrams under topological isotopy. It is significant that different algebraic conditions have the same topological representation. That means that diagrams that are equivalent up to isotopy have different meanings in quantum information when they are located at different positions. In our two-string model we have used this same philosophy to design protocols in a topological way [6, 7].

Using these diagrams, we obtain a 3D representation of the CNOT map and the teleportation protocol. The teleportation protocol becomes a topological protocol in the quon model.

## 2. PARAFERMION ALGEBRAS

The *parafermion algebra*  $PF_n$  of order  $d$  is a  $*$ -algebra with unitary generators  $c_m$ ,  $m = 1, 2, \dots, n$ , which satisfy

$$c_m^d = 1 \quad \text{and} \quad c_m c_{m'} = q c_{m'} c_m \quad \text{for} \quad 1 \leq m < m' \leq n. \quad (1)$$

Here  $q \equiv e^{\frac{2\pi i}{d}}$  and  $i \equiv \sqrt{-1}$ . Consequently  $c_j^* = c_j^{-1} = c_j^{d-1}$ , where  $*$  denotes the adjoint. Majorana fermions arise from the case  $d = 2$ . The Jordan-Wigner transformation gives the isomorphism  $PF_{2n} \cong M_d(\mathbb{C})^{\otimes n}$  to tensor products of  $d \times d$  matrices. The parafermion algebra  $PF_n$  has a basis  $\{c^\alpha\}$ . Here  $c^\alpha = c_1^{\alpha_1} \dots c_n^{\alpha_n}$ , and  $\alpha_k \in \mathbb{Z}_d$ . The charge of  $c^\alpha$  is defined to be  $|c^\alpha| = \sum_{k=1}^n \alpha_k$  in  $\mathbb{Z}_d$ . The zero-charged elements  $PF_n^0$  form a subalgebra of the parafermion algebra  $PF_n$ , namely the neutral subalgebra.

**2.1. Parafermion planar para algebras (PAPPA).** Given  $1 \leq m \leq n$ , our diagrammatic representation for  $c_m^{\alpha_m}$ , with  $\alpha_m = k$ , as explained in [3], is

$$c_m^k \longleftrightarrow \left| \dots \left| k \right| \dots \right|,$$

where the  $m^{\text{th}}$  string is labeled by  $k$ . In our notation we have the relations

$$\begin{aligned} \textbf{Multiplication:} \quad & \left| \begin{array}{c} \ell \\ k \end{array} \right| = \left| \begin{array}{c} k + \ell \\ k \end{array} \right|, \quad d \left| \begin{array}{c} \ell \\ \ell \end{array} \right| = \left| \begin{array}{c} 0 \\ 0 \end{array} \right|. \\ \textbf{Para isotopy:} \quad & \left| \begin{array}{c} \ell \\ k \end{array} \right| \left| \dots \right| \left| \begin{array}{c} \ell \\ k \end{array} \right| = q^{k\ell} \left| \begin{array}{c} k \\ k \end{array} \right| \left| \dots \right| \left| \begin{array}{c} \ell \\ \ell \end{array} \right|. \end{aligned} \quad (2)$$

The strings between  $k$ -charged and  $\ell$ -charged strings are not charged. We call  $q^{k\ell}$  the twisting scalar.

Take  $\zeta$  to be a square root of  $q$ , such that  $\zeta^{d^2} = 1$ . We can interpolate between the diagrams in (2) as a twisted tensor product:

$$\textbf{Twisted product:} \quad \left| \begin{array}{c} \ell \\ k \end{array} \right| \left| \dots \right| \left| \begin{array}{c} \ell \\ \ell \end{array} \right| = \zeta^{k\ell} \left| \begin{array}{c} k \\ k \end{array} \right| \left| \dots \right| \left| \begin{array}{c} \ell \\ \ell \end{array} \right|. \quad (3)$$

In the PAPPA model, the charged strings on the 2D plane satisfy the following relations:

- $\rho_\pi(c^j) = \zeta^{j^2} c^j$ , where  $\rho_\pi$  is a rotation by  $\pi$  on the plane. Then  $\rho_{2\pi}(c^j) = q^{j^2} c^j$  for the  $2\pi$  rotation  $\rho_{2\pi}$ .
- $j \bigcirc = 0$ ,  $1 \leq j \leq d-1$ , and  $\bigcirc = \delta = \sqrt{d}$ .

Moreover, we can introduce additional relations:

- Let  $\omega = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} \zeta^{j^2}$ . This is a phase, as shown in Proposition 2.15 of [3]. Then

$$\begin{aligned} \left| \begin{array}{c} \ell \\ k \end{array} \right| \left| \begin{array}{c} \ell \\ k \end{array} \right| &= \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} \left( \begin{array}{c} -j \\ j \end{array} \right) \\ \times &= \frac{1}{\sqrt{\omega d}} \sum_{j=0}^{d-1} \zeta^{j^2} \left( \begin{array}{c} -j \\ j \end{array} \right). \end{aligned}$$

Using the above definition of braiding, we can establish the braided relation that any neutral diagram can move above or under the strings, see [6]. Therefore the neutral diagrams can be lifted to the 3D space.

**2.2. Categorical approach to the neutral part of the parafermion algebra.** For readers who are familiar with category theory, one can consider the neutral diagrams as morphisms in a monoidal category. The neutral part of PAPP is the  $\mathbb{Z}_d$  unshaded subfactor planar algebra. It is a  $\mathbb{Z}_2$  graded unitary fusion category. Its even part is the monoidal category  $Vec_{\mathbb{Z}_d}$ , whose simple objects  $X_g$  are labeled by ground elements  $g$  in  $\mathbb{Z}_d$  indicating the fusion rule. It has only one odd simple object  $\tau = \bar{\tau}$ , such that  $\tau^2 = \gamma = \bigoplus_{g \in \mathbb{Z}_d} X_g$ . Thus  $\gamma$  is a Frobenius algebra. Then the neutral subalgebra of the parafermion algebra is given by  $PF_n^0 = \text{hom}(\tau^n, \tau^n)$ .

### 3. THE QUON MODEL

The neutral diagrams with four boundary points are  $d$ -dimensional. We use these diagrams to represent a qudit vector as described in §5.3 of [3]. The general framework that we introduce is to embed the neutral qudits into a three manifold.

**3.1. Qudits.** An  $n$ -qudit is represented by  $n$  hemispheres, such that each hemisphere contains a neutral diagram with four boundary points on its boundary disc. For example, the 3-qudit is represented as

$$(4)$$

Here  $v_j$  labels a 1-qudit vector given by neutral diagrams with four boundary points in the hemisphere. Here we allocate the boundary of the hemispheres on the  $X$ - $Y$ -plane in the 3D space.

**3.2. Transformations.** An  $n$ -qudit transformation is represented by a neutral element  $T$  in  $PF_{4n}^0 = \text{hom}(\tau^{4n}, \tau^{4n})$  in a 3-manifold. The 3-manifold has  $n$  boundary discs on the top and  $n$  at the bottom. Each disc contains four boundary points of  $T$ . For example, a 3-qudit transformation  $T$  will have the representation

$$(5)$$

**3.3. Isotopy of Neutral Diagrams.** In addition to the relations for charged strings, we allow isotopy of strings in 3-manifolds. We define a relation for 3-manifolds: if a 3-sphere has no diagram inside, then it can be removed. Moreover we define a joint relation between diagrams and 3-manifolds.

Suppose  $T \in \text{hom}(\tau^m, \tau^n)$ . Let  $B_m$  be an orthonormal basis (ONB) of  $\text{hom}(\tau^m, \tau^m)$ , and let  $B_n$  be an ONB of  $\text{hom}(\tau^n, \tau^n)$ . We define the relation

$$(6)$$


Note that the second equality always holds. The proof is from basic linear algebra. It tells that the relation is well-defined up to isotopy of neutral diagrams in 3-manifolds. By this relation, the picture (5) reduces to a




These are the three eigenbases of the three unitary Pauli matrices. The matrices are given diagrammatically in [3], respectively as

$$Z = \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline -1 \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array}, \quad X = \begin{array}{|c|} \hline 1 \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline -1 \\ \hline \end{array}, \quad Y = \begin{array}{|c|} \hline -1 \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline 1 \\ \hline \end{array}. \quad (8)$$

In the quon model we represent the basis of the 1-qudit space by a pair of strings with opposite charges, embedded in a hemisphere:



The algebraic adjoint operation is given by a geometric reflection along the  $Z$ -direction. Therefore a measurement is represented by a pair of strings with opposite charges in the reflected hemisphere:



The charge represents the result of the measurement.

### 5. 1-QUDIT CLIFFORD GROUP

The 1-qudit transformations  $\{X, Y, Z, F, G\}$  are generators of the 1-qudit Clifford group. Their algebraic definitions are given by:

$$\begin{aligned} X|k\rangle &= |k+1\rangle, \quad Y|k\rangle = \zeta^{1-2k}|k-1\rangle, \quad Z|k\rangle = q^k|k\rangle, \\ F|k\rangle &= \frac{1}{\sqrt{N}} \sum_{i=0}^{N-1} q^{ki} |i\rangle, \quad G|k\rangle = \zeta^{k^2}|k\rangle. \end{aligned} \quad (9)$$


When we consider 1-qudit as a vector state, the qudit transformations are defined up to a phase. Thus 1-qudit transformations form a group  $\mathbb{Z}_d^2 \times SL(2, \mathbb{Z}_d)$  as shown in [3]. The Pauli matrices are given diagrammatically by (8), while

$$F = \begin{array}{|c|} \hline \diagdown \\ \hline \end{array} \begin{array}{|c|} \hline \diagup \\ \hline \end{array} = \begin{array}{|c|} \hline \diagup \\ \hline \end{array} \begin{array}{|c|} \hline \diagdown \\ \hline \end{array}, \quad G = \begin{array}{|c|} \hline \diagdown \\ \hline \end{array} \begin{array}{|c|} \hline | \\ \hline \end{array} = \begin{array}{|c|} \hline | \\ \hline \end{array} \begin{array}{|c|} \hline \diagdown \\ \hline \end{array}. \quad (10)$$


### 6. N-QUDIT CLIFFORD GROUP

Let  $C_X$  be the controlled  $X$  transformation. For the qubit case it becomes CNOT. The  $n$ -qudit clifford group is generated by  $\{X, Y, Z, F, G, C_X\}$ . We represent the qudit transformation  $C_X$  by neutral diagrams in 3-manifolds. The projection of neutral diagrams on the 2D plane is represented as follows. It turns out to be more natural to draw the picture in the 3D space. We discuss more about  $C_X$  and the 3D representations in §9–11.

2D projection of CNOT



3D CNOT



7. RESOURCE STATES


The generalized Bell state for qudits is given by  $d^{-1/2} \sum_{k \in \mathbb{Z}_d} |k, k\rangle$ . Its diagrammatic representation is



One can check this identification by the joint relation.

8. TELEPORTATION

In the quon model, the teleportation protocol is represented by the following diagrammatic protocol:



The pair of opposite-charged-strings is neutral, thus it is defined in the 3D space. It obviously represents the teleportation process in a topological way. Moreover, it shows the one-to-one correspondence between the diagrammatic representation of this protocol and the algebraic representation in the teleportation protocol of Bennett et al. [17].

9. BI-FROBENIUS ALGEBRAS

In tensor networks, the CNOT gate on qubits is decomposed as two transformations COPY and SUM. They are defined as follows algebraically and represented by planar diagrams:

$$\begin{aligned}
 CNOT &= \sum_{k, j \in \mathbb{Z}_d} |k + j, j\rangle \langle k, j| : \begin{array}{c} | \\ \bullet \\ | \end{array} \begin{array}{c} | \\ \circ \\ | \end{array} \\
 COPY &= \sum_{j \in \mathbb{Z}_d} |j, j\rangle \langle j| : \begin{array}{c} | \\ \bullet \\ / \end{array} \begin{array}{c} | \\ \circ \\ | \end{array} \\
 SUM &= \sum_{j \in \mathbb{Z}_d} |k + j\rangle \langle k, j| : \begin{array}{c} | \\ \bullet \\ / \end{array} \begin{array}{c} | \\ \circ \\ \backslash \end{array}
 \end{aligned}$$



In the quon model, we can represent these maps in a consistent way by strings in 3-manifolds, such that the algebraic relations become topological isotopy. The COPY map is represented by



One can check that this diagrammatic definition coincides with the algebraic definition by (6). By this representation, the algebraic conditions of the Frobenius algebra become topological isotopy in the quon model.

### 10. THE STRING-GENUS JOINT RELATION


In case we have a diagram of the form: a closed neutral string surrounds a genus of the manifold, then we can remove both, up to a scalar. When  $m$  and  $n$  are odd numbers, we have

Note that  $\tau^m, \tau^n$  are multiples of  $\tau$ . It is enough to prove the relation for  $m = n = 1$ . In this case, it follows from the relation (6).

If  $m$  or  $n$  is even, by relation (6), then the diagram is 0. Thus if the diagram is a part of a non-zero transformation, then the  $m$  and  $n$  have to be odd numbers. In this case, the relation means that if there is a circle around a genus of the 3-manifold, then we can remove the circle and the genus by multiplying a scalar  $d^{-1/2}$ .

### 11. TOPOLOGICAL RELATIONS FOR $C^*$ -HOPF ALGEBRAS

A more conceptual way to look at picture in the quon model is to assign the four boundary points of the quon to the corners of a square in a plane orthogonal to the  $Z$ -axis. The white and black bullets indicate diagrammatic operations in  $X$  and  $Y$ -directions on the 2D plane. For example, if we look at this 3D diagram from the top along the  $Z$ -direction, then the pictures for COPY and SUM become respectively



Here we only draw the boundary circle of the 3-manifold to simplify the picture. Similarly we can represent the white and black spiders as strings in 3-manifolds in the quon model. The white spiders are expended in the  $X$ -direction and the black spiders are expended in the  $Y$ -direction. From this point of view, the Fourier transform is a  $90^\circ$  rotation around the  $Z$ -axis, which explains the duality of the two Frobenius algebras in

(12) in a geometric way. Moreover, one can check that the relation (15) also becomes an isotopy in the 3D space in the quon model.

Furthermore, this pair of Frobenius algebras satisfies the following additional relations (17)–(20). These relations define  $\mathbb{Z}_d$  as a  $C^*$  Hopf algebra, where  $F^2$  is the antipode map of the Hopf algebra and the involution is an anti-linear map which reflects the diagrams vertically. Note that if a pair of Frobenius algebras satisfy these relations, then the underlying  $d$  dimensional Hilbert space becomes a Hopf algebra. This has been observed in [11].

$$\text{Diagram (17)} = \text{Diagram (17)}$$

$$\text{Diagram (18)} = \text{Diagram (18)}$$

$$\text{Diagram (19)} = \text{Diagram (19)} = \text{Diagram (19)}$$

$$\text{Diagram (20)} = \text{Diagram (20)}$$

(17)

(18)

(19)

(20)

Now we give an interpretation of these relations as topological isotopy in 3D. Note that relation (17) follows from the definition of the COPY map. By isotopy in our quon model, relations (18) and (19) are exactly the same diagrams as the relation (17). The most interesting relation is (20), and we explain that relation in detail.

Relation (20) becomes topological isotopy, when we use the string-genus joint relation established in §10, namely the joint relation (16). Then (20) is given by the isotopy:

$$\text{Diagram} = d^{-1/2} \text{Diagram}$$

Thus we have given a topological interpretation for the  $C^*$  Hopf algebra axioms for  $\mathbb{Z}_d$ .

There is a one-to-one correspondence between  $C^*$  Hopf algebras and irreducible, depth-two subfactor planar algebras [18, 19]. In this case,  $d$  is the global dimension of the  $C^*$ -Hopf algebra. Moreover, the even part of the planar algebra is the representation category of the Kac algebra. The odd part has only one simple object  $\tau$ , so (16) also holds. In the above interpretation, we only use (shaded) planar diagrams without braids in 3-manifolds. So this topological interpretation works for any finite dimensional  $C^*$  Hopf algebra. From this point of view, many algebraic properties of  $C^*$ -Hopf algebras reduce to topological isotopy.

## 12. QUON MODEL OF A UNITARY MODULAR TENSOR CATEGORY

We can define the quon model for any unitary modular tensor category  $\mathcal{C}$ , so that the 1-quon basis corresponds to the set of simple objects  $OB$  in  $\mathcal{C}$ . If we take  $\mathcal{C}$  to be the unitary modular tensor category, such that its fusion ring is  $Z_d$  and its modular  $S$  matrix is  $q^{kl}$ , where  $q = e^{\frac{2\pi i}{d}}$ , then we get back the quon model for qudits defined by PAPPÀ.

For each  $X \in OB$ , we obtain a simple object  $\tilde{X} := X \otimes \bar{X}$  in  $\mathcal{C} \otimes \mathcal{C}$ , where  $\bar{X}$  is the dual of  $X$ . Take  $\gamma = \bigoplus_{X \in OB} X \otimes \bar{X}$  in  $\mathcal{C} \otimes \mathcal{C}$ . It is known that  $\gamma = \bigoplus_{X \in Ob} X \otimes \bar{X}$  is a Frobenius algebra in  $\mathcal{C} \otimes \mathcal{C}$ . Thus  $\mathcal{P}_{n,+} = \text{hom}(1, \gamma^n)$  is a subfactor planar algebra generated by  $\tau$ , such that  $\gamma = \tau \otimes \bar{\tau}$ . It is proved in [20] that this planar algebra is unshaded. That means  $\tau = \bar{\tau}$ .

Note that  $\text{hom}(1, \tau^4) \cong \text{hom}(1, \gamma^2)$ . This space has an orthonormal-basis given by the canonical inclusion from 1 to  $(X \otimes \bar{X}) \otimes (\bar{X} \otimes X)$ , for  $X \in OB$ . We call the generalized single-particle state a *1-quon*. Thus the dimension  $d$  of the 1-quon space is the cardinality of  $OB$ . Furthermore, it is proved in [20] that the string Fourier transform on the 1-quon space is the  $S$  matrix of the unitary modular tensor category  $\mathcal{C}$ .

We generalize the quon model as follows: We label each of the four boundary points by the object  $\tau$ . The diagrams in the 3-manifolds are given by morphisms in  $\mathcal{C} \otimes \mathcal{C}$ . The representation for a  $n$ -quon is given by morphisms in  $\text{hom}(1, \tau^4)$  in  $n$  hemispheres as in (4). The  $n$ -quon transformations are represented by morphisms in  $\text{hom}(\tau^{4n}, \tau^{4n})$  in a 3-manifold as in (5). The relations between diagrams and 3-manifolds are also defined by (6). Then  $n$ -quon transformations also reduce to a linear sum of the form in (7) which represent matrix units. Therefore the  $n$ -quon transformations are transformations on the  $d^n$  dimensional Hilbert space.

In general, the quon model can be defined for any subfactor planar algebra [21], if we do not require  $\tau = \bar{\tau}$ . In this case, the diagrams in the 3-manifolds with  $4n$  boundary points are given by a shaded planar diagram in the  $2n$ -box space of planar algebras. We have used this general case to give the topological interpretation of the  $C^*$ -Hopf algebra relations.

## 13. ACKNOWLEDGEMENT

We are grateful to acknowledge an interesting discussion with Daniel Loss about his paper [22], and with Michael Freedman about his paper [23]. We thank Bob Coecke for sharing a manuscript of the book [11]. This research was supported by a grant from the Templeton Religion Trust.

## REFERENCES

- [1] A. Kitaev, Fault-tolerant quantum computation by anyons, *Ann. Phys.* **303** (2003), 2–30, <https://arxiv.org/abs/quant-ph/9707021>, doi:10.1016/S0003-4916(02)00018-0.
- [2] M. H. Freedman, A. Kitaev, M. J. Larsen, and Z. Wang, Topological quantum computation, *Bulletin of the American Mathematical Society* Volume 40, Number 1, (2002), 31–38, doi:10.1090/S0273-0979-02-00964-3.
- [3] A. M. Jaffe and Z. Liu, Planar para algebras, reflection positivity, *Commun. Math. Phys.* to appear, <http://arxiv.org/abs/1602.02662>.
- [4] A. M. Jaffe, Z. Liu, and A. Wozniakowski, Qudit isotopy, <https://arxiv.org/abs/1602.02671>.
- [5] A. Kitaev, Anyons in an exactly solved model and beyond, *Ann. Phys.* **306** (2006) 2–111, <http://dx.doi.org/10.1016/j.aop.2005.10.005>.
- [6] A. M. Jaffe, Z. Liu, and A. Wozniakowski, Holographic software for quantum networks, preprint, <http://arxiv.org/abs/1605.00127>.
- [7] A. M. Jaffe, Z. Liu, and A. Wozniakowski, Topological design of protocols, <https://arxiv.org/abs/1605.00321>.
- [8] S. Abramsky and B. Coecke, A categorical semantics of quantum protocols, *Logic in Computer Science, 2004. Proceedings of the 19th Annual IEEE Symposium* IEEE (2004).
- [9] B. Coecke and R. Duncan, Interacting quantum observables. In *Automata, Languages and Programming*, **5126** 298–310 (2008), Springer, doi:10.1007/978-3-540-70583-3\_25.

- [10] B. Coecke and R. Duncan, Interacting quantum observables: categorical algebra and diagrammatics, *New Journal of Physics*, **13**(4) 043016 (2011), [doi:10.1088/1367-2630/13/4/043016](https://doi.org/10.1088/1367-2630/13/4/043016).
- [11] B. Coecke and A. Kissinger, Picturing quantum processes: A first course in quantum theory and diagrammatic reasoning, Cambridge University Press, to appear 2017, ISBN:9781107104228.
- [12] V. Bergholm and J. Biamonte, Categorical quantum circuits, *J. Phys A: Math. Theor.* **44** 245304 (2011), <https://arxiv.org/abs/1010.4840>.
- [13] S. J. Denny, J. D. Biamonte, D. Jaksch, and S. R. Clark Algebraically contractible topological tensor network states, *J. Phys. A: Math. Theor.* **45** 015309 (2012), <https://arxiv.org/abs/1108.0888>.
- [14] M. K. Backens, Completeness and the ZX-calculus, Thesis, Oxford University, <https://arxiv.org/abs/1602.08954>.
- [15] J. Vicary, Higher quantum theory, <https://arxiv.org/abs/1207.4563>.
- [16] D. Reutter and J. Vicary, Biunitary constructions in quantum information, <https://arxiv.org/abs/1609.07775>.
- [17] C. H. Bennett, G. Brassard, C. Crépeau, R. Jozsa, A. Peres, and W. K. Wootters, Teleporting an unknown quantum state via dual classical and Einstein-Podolsky-Rosen Channels, *Phys. Rev. Lett.* **70**, (1993) 1895, [doi:10.1103/PhysRevLett.70.1895](https://doi.org/10.1103/PhysRevLett.70.1895).
- [18] W. Szymanski, Finite index subfactors and Hopf algebra crossed products *Proc. Amer. Math. Soc.* **120**, (1994) no. 2, 519-528.
- [19] V. Kodyalam, Z. Landau, and V. S. Sunder, The planar algebra associated to a Kac algebra, *Proceedings of the Indian Academy of Sciences-Mathematical Sciences.* (2003) Vol. 113. No. 1. Springer India.
- [20] Z. Liu, and F. Xu, Jones-Wassermann subfactors for modular tensor categories, <https://www.arxiv.org/abs/1612.xxxx>.
- [21] V. F. R. Jones, Planar algebras I., *New Zealand Journal of Mathematics* (1998), <https://arxiv.org/abs/math/9909027>.
- [22] A. Hutter and D. Loss, Quantum computing with parafermions, *Phys. Rev.* **B 93**, (2016) 125105 1–7, [doi:10.1103/PhysRevB.93.125105](https://doi.org/10.1103/PhysRevB.93.125105).
- [23] T. Karzig, C. Knapp, R. M. Lutchyn, P. Bonderson, M. Hastings, C. Nayak, J. Alicea, K. Flensberg, S. Plugge, Y. Oreg, C. Marcus, and M. H. Freedman, Scalable designs for quasiparticle-poisoning-protected topological quantum computation via Majorana zero modes, <https://arxiv.org/abs/1610.05289>.  
*E-mail address:* zhengweiliu@fas.harvard.edu  
  
*E-mail address:* airwozz@gmail.com  
  
*E-mail address:* arthur\_jaffe@harvard.edu